

An $\mathcal{O}(h^8)$ optimal B-spline collocation for solving higher order boundary value problems

Shokofeh Sharifi[†] and Jalil Rashidinia^{‡*}

[†]*Department of Mathematics and statistics, Central Tehran Branch,
Islamic Azad University, Tehran, Iran*

[‡]*School of Mathematics, Iran University of Science and Technology,
Tehran, Iran*

Emails: sh.sharifi_m61@yahoo.com, rashidinia@iust.ac.ir

Abstract. As we know the approximation solution of seventh order two points boundary value problems based on B-spline of degree eight has only $\mathcal{O}(h^2)$ accuracy and this approximation is non-optimal. In this work, we obtain an optimal spline collocation method for solving the general nonlinear seventh order two points boundary value problems. The $\mathcal{O}(h^8)$ convergence analysis, mainly based on the Green's function approach, has been proved. Numerical illustration demonstrate the applicability of the purposed method. Three test problems have been solved and the computed results have been compared with the results obtained by recent existing methods to verify the accurate nature of our method.

Keywords: Nonlinear boundary value problems, eighth degree B-spline, collocation method, convergence analysis, Green's function.

AMS Subject Classification: 65L10, 65L12, 65L20, 65L70.

1 Introduction

We consider the general nonlinear seventh order two point boundary value problems (BVPs) of the following form:

$$Ly \equiv y^{(7)}(x) - f(x, y(x), y'(x), \dots, y^{(6)}(x)) = 0, \quad a \leq x \leq b, \quad (1)$$

*Corresponding author.

with the boundary conditions,

$$By \equiv \sum_{j=0}^6 (\alpha_{ij} y^{(j)}(a) + \beta_{ij} y^{(j)}(b)) = \eta_i, \quad 0 \leq i \leq 6, \quad (2)$$

where α_{ij} , β_{ij} and η_i are given real constants, f is a continuous function, $y(x)$ is an unknown function, and L and B are differential operators.

The formulation of many mathematical models in engineering and other branches of sciences are in the form of differential equations with initial or boundary conditions and boundary value problems generally. Obtaining the analytic solution for these problems is impossible, because of this, many authors attempt to use different numerical methods such as finite difference, Galerkin and Sinc collocation methods [12, 14, 15].

The literature on the numerical solution of seventh order two point boundary value problems is seldom. These problems are generally arise in modelling induction motors with two rator circuits. Behaviour of such models have been studied by Richards and Sarma [18]. The solution of seventh order BVPs based on variational iteration and differential transformation method are given by Siddiqi et al. [21, 22]. In [23] the authors used the homotopy analysis method for solving higher order BVPs. Reproducing kernel method for the solution of seventh order BVPs has been studied in [3].

Many researchers applied the collocation methods for solution of BVPs [2, 4, 19]. The spline functions has been applied to solve BVPs in [5], with order $\mathcal{O}(h^2)$. After that many authors [1, 7, 8, 20] examined the collocation method based on cubic spline for BVPs.

An optimal cubic spline collocation method at grid points was developed by Danial and Swatz in [6], which gives $\mathcal{O}(h^4)$ accuracy. In [10] the authors used optimal collocation method on midpoints based on quadratic spline for approximate the solution of second order BVPs. Irodou-Ellina and Houstis applied the optimal quintic spline collocation method for solving linear fourth order two point BVPs which lead to an $\mathcal{O}(h^6)$ approximation [11].

In [16] Rashidinia et al. developed an optimal method based on sextic spline at the grid points for solving of nonlinear fifth order two point BVPs. Also, Rashidinia and Ghasemi [17] applied an optimal sextic spline at the midpoints for the numerical solution of sixth order nonlinear two point BVPs.

In this paper we applied optimal collocation method based on B-spline of degree eight at the nodal points of the interval $[a, b]$ and obtained $\mathcal{O}(h^8)$ approximation for the numerical solution of boundary value problems (1) – (2). The approximation is assumed to satisfy a high order approximation

of the problem. In Section 2, we obtain the consistency relations for spline of degree eight at the nodal points of the partition. In Section 3, the description of the method based on spline for the solution of (1) – (2) is explained. The convergence analysis of the presented method is given in detail, in Section 4. In Section 5, numerical experiments are conducted to demonstrate the applicability of the proposed method computationally. Conclusion is presented in Section 6.

2 Spline interpolation

We define the spline of degree eight as basis functions to construct an interpolant $S(x)$, satisfying certain end conditions and then derive several relations that are useful in the formulation of the optimal spline collocation method.

Now let $\Delta \equiv \{a = x_0 < x_1 < \dots < x_n = b\}$ be a uniform partition of the interval $[a, b]$ with the step size $h = \frac{b-a}{n}$. We consider smooth spline of degree eight $S(x)$, that is an element of $Sp_8(\Delta) \equiv \{q(x) | q(x) \in C^7[a, b]\}$ and $q(x)$ is a polynomial of degree at most 8 on the partition Δ . The set of B-splines $\{B_k(x)\}_{k=-3}^{n+4}$, form a basis for $Sp_8(\Delta)$, so we can define our spline of degree eight in the following form:

$$S(x) = \sum_{k=-3}^{n+4} c_k B_k(x), \quad x \in [x_i, x_{i+1}],$$

that satisfies the following interpolatory conditions:

$$S(x_i) = y(x_i), \quad 0 \leq i \leq n, \quad (3)$$

associated with the end conditions:

$$S^{(7)}(x_i) = y^{(7)}(x_i) - \frac{h^2}{12} y^{(9)}(x_i) + \frac{h^4}{240} y^{(11)}(x_i) - \frac{h^6}{6048} y^{(13)}(x_i), \quad (4)$$

for $i = 0, 1, 2, 3, n-2, n-1, n$. By using linear dependence relations, we have the following consistency relations for spline of degree eight and its first seventh derivatives for $4 \leq i \leq n-3$ at the grid points: [9, 24]

$$\begin{aligned} (a) \quad \Gamma S_i^{(7)} &= \frac{40320}{h^7} (\mp S_{i-4, i+3} \pm 7S_{i-3, i+2} \mp 21S_{i-2, i+1} \pm 35S_{i-1, i}), \\ (b) \quad \Gamma S_i^{(6)} &= \frac{20160}{h^6} (S_{i-4, i+3} - 5S_{i-3, i+2} + 9S_{i-2, i+1} - 5S_{i-1, i}), \\ (c) \quad \Gamma S_i^{(5)} &= \frac{6720}{h^5} (\mp S_{i-4, i+3} \pm S_{i-3, i+2} \pm 9S_{i-2, i+1} \mp 25S_{i-1, i}), \\ (d) \quad \Gamma S_i^{(4)} &= \frac{1680}{h^4} (S_{i-4, i+3} + 7S_{i-3, i+2} - 27S_{i-2, i+1} + 19S_{i-1, i}), \end{aligned}$$

$$\begin{aligned}
(e) \quad \Gamma S_i^{(3)} &= \frac{336}{h^3} (\mp S_{i-4,i+3} \mp 23S_{i-3,i+2} \pm 9S_{i-2,i+1} \pm 95S_{i-1,i}), \\
(f) \quad \Gamma S_i^{(2)} &= \frac{56}{h^2} (S_{i-4,i+3} + 55S_{i-3,i+2} + 189S_{i-2,i+1} - 245S_{i-1,i}), \\
(g) \quad \Gamma S_i^{(1)} &= \frac{8}{h} (\mp S_{i-4,i+3} \mp 119S_{i-3,i+2} \mp 1071S_{i-2,i+1} \mp 1225S_{i-1,i}), \\
(h) \quad \Gamma g_i &= (g_{i-4,i+3} + 247g_{i-3,i+2} + 4293g_{i-2,i+1} + 15619g_{i-1,i}), \quad (5)
\end{aligned}$$

where the discrete operator Γ is defined for any function g on the interval $[a, b]$. For sake of convenience we set $S_i \equiv S(x_i)$, $S_i^j \equiv S^j(x_i)$, $i = 0, 1, \dots, n$, $j = 1, 2, \dots, 7$ where $g^{(j)} \equiv D^{(j)}g$. In order to obtain the error bounds for spline of degree eight S and its derivatives $S', \dots, S^{(7)}$, we present the next theorem.

Theorem 1. *Let $S(x)$ be the spline of degree eight, satisfying (3) – (4) and interpolating the function $y \in C^{14}[a, b]$, then for $i = 0, 1, \dots, n$ the following relations hold,*

$$\begin{aligned}
(a) \quad S_i^{(1)} &= y_i^{(1)} + \mathcal{O}(h^8), \\
(b) \quad S_i^{(2)} &= y_i^{(2)} + \mathcal{O}(h^8), \\
(c) \quad S_i^{(3)} &= y_i^{(3)} - \frac{h^6}{30240}y_i^{(9)} + \mathcal{O}(h^8), \\
(d) \quad S_i^{(4)} &= y_i^{(4)} + \frac{h^6}{6048}y_i^{(10)} + \mathcal{O}(h^8), \\
(e) \quad S_i^{(5)} &= y_i^{(5)} + \frac{h^4}{720}y_i^{(9)} - \frac{h^6}{3024}y_i^{(11)} + \mathcal{O}(h^8), \\
(f) \quad S_i^{(6)} &= y_i^{(6)} - \frac{h^4}{240}y_i^{(10)} + \frac{h^6}{3024}y_i^{(12)} + \mathcal{O}(h^8), \\
(g) \quad S_i^{(7)} &= y_i^{(7)} - \frac{h^2}{12}y_i^{(9)} + \frac{h^4}{240}y_i^{(11)} - \frac{h^6}{6048}y_i^{(13)} + \mathcal{O}(h^8), \quad (6)
\end{aligned}$$

and we have the following error bounds,

$$\| (S - y)^{(k)} \| = \mathcal{O}(h^{9-k}), \quad k = 1, 2, \dots, 7. \quad (7)$$

Proof. First we need to prove relation (6g). Using Taylor's series expansion and taking into account the interpolatory condition $S_i = y_i$, $i = 0, 1, \dots, n$, in the relation (5a) we have

$$\begin{aligned}
\Gamma S_i^{(7)} &= 40320y_i^{(7)} - 20160hy_i^{(8)} + 16800h^2y_i^{(9)} - 6720h^3y_i^{(10)} + 3192h^4y_i^{(11)} \\
&\quad - 1064h^5y_i^{(12)} + \frac{1120}{3}h^6y_i^{(13)} - \frac{320}{3}h^7y_i^{(14)} + \mathcal{O}(h^8), \quad (8)
\end{aligned}$$

for $4 \leq i \leq n-3$. Further, using Taylor's series expansion, for any function $g \in C^8[a, b]$ we obtain

$$\begin{aligned} \Gamma g_i = & 40320g_i - 20160hg'_i + 20160h^2g''_i - 8400h^3g'''_i + 4704h^4g^{(4)}_i \\ & - 1680h^5g^{(5)}_i + 688h^6g^{(6)}_i - 215h^7g^{(7)}_i + \mathcal{O}(h^8), \quad 4 \leq i \leq n-3. \end{aligned} \quad (9)$$

Setting $g(x) = y_i^{(7)} - \frac{h^2}{12}y_i^{(9)} + \frac{h^4}{240}y_i^{(11)} - \frac{h^6}{6048}y_i^{(13)}$, we have the following relation

$$\begin{aligned} \Gamma g_i = & \Gamma(y_i^{(7)} - \frac{h^2}{12}y_i^{(9)} + \frac{h^4}{240}y_i^{(11)} - \frac{h^6}{6048}y_i^{(13)}) \\ = & 40320y_i^{(7)} - 20160hy_i^{(8)} + 16800h^2y_i^{(9)} - 6720h^3y_i^{(10)} + 3192h^4y_i^{(11)} \\ = & -1064h^5y_i^{(12)} + \frac{1120}{3}h^6y_i^{(13)} - \frac{320}{3}h^7y_i^{(14)} + \mathcal{O}(h^8). \end{aligned} \quad (10)$$

By subtracting Eq. (8) from (10), we obtain

$$\Gamma(S_i^{(7)} - y_i^{(7)} + \frac{h^2}{12}y_i^{(9)} - \frac{h^4}{240}y_i^{(11)} + \frac{h^6}{6048}y_i^{(13)}) = \mathcal{O}(h^8), \quad 4 \leq i \leq n-3. \quad (11)$$

Denoting $R_i \equiv S_i^{(7)} - y_i^{(7)} + \frac{h^2}{12}y_i^{(9)} - \frac{h^4}{240}y_i^{(11)} + \frac{h^6}{6048}y_i^{(13)}$, then by associating the Eq. (4) and consistency equation (11), we get the following system of equations,

$$\Gamma R_i = \mathcal{O}(h^8) \quad \forall y^{(14)}, \quad 4 \leq i \leq n-3,$$

$$R_0 = R_1 = R_2 = R_3 = R_{n-2} = R_{n-1} = R_n = 0. \quad (12)$$

Since the coefficient matrix of the above system is positive definite, it is nonsingular and its inverse has a finite norm. Thus we have $R_i = \mathcal{O}(h^8)$, $i = 0, 1, \dots, n$, this concludes the proof of relation (6g).

To prove relation (6b) consider the following relations, which can be easily obtained via long straightforward calculations for any spline of degree eight at the interior grid points x_i ,

$$\begin{aligned} S_i^{(6)} = & \frac{-1}{40320h^6}[-40320S_{i,i+6} + 241920S_{i+1,i+5} - 604800S_{i+2,i+4} \\ & + 806400S_{i+3} - h^7(20159S_i^{(7)} + 40072S_{i+1}^{(7)} + 35779S_{i+2}^{(7)} + 20160S_{i+3}^{(7)} \\ & + 4541S_{i+4}^{(7)} + 248S_{i+5}^{(7)} + S_{i+6}^{(7)})], \quad 0 \leq i \leq n-6, \end{aligned}$$

and

$$S_i^{(6)} = \frac{-1}{40320h^6} [\pm 116081280S_{i-7,i} \mp 812609280S_{i-6,i-1} \pm 2437948800S_{i-5,i-2} \\ \mp 4063449600S_{i-4,i-3} + h^7(2879S_{i-7}^{(7)} + 711112S_{i-6}^{(7)} + 12359299S_{i-5}^{(7)} \\ + 44962560S_{i-4}^{(7)} + 44946941S_{i-3}^{(7)} + 12323768S_{i-2}^{(7)} + 671041S_{i-1}^{(7)} \\ - 17280S_i^{(7)})], \quad 7 \leq i \leq n.$$

Using relation part (g) of Eq. (6) in the above relations and applying Taylor's series expansion of $y_{i\pm l}^{(k)}$ for $k = 0, 7, 9, 11$ we get

$$S_i^{(6)} = y_i^{(6)} - \frac{h^4}{240}y_i^{(10)} + \frac{h^6}{3024}y_i^{(12)} + \mathcal{O}(h^8), \quad 0 \leq i \leq n.$$

In a similar manner applying some appropriate consistency relations we can prove the other relations in this Theorem. \square

To improve the order of numerical solution of the system of equations (1)–(2) we need to donate the following discrete operators for convenience:

$$\begin{aligned} \lambda_0 g_i &= g_{i-3,i+3} - 6g_{i-2,i+2} + 15g_{i-1,i+2} - 20g_i, \quad 3 \leq i \leq n-3 \\ \lambda_1 g_i &= -\frac{1}{6}[g_{i-3,i+3} - 12g_{i-2,i+2} + 39g_{i-1,i+1} - 56g_i], \quad 3 \leq i \leq n-3 \\ \lambda_2 g_i &= -\frac{1}{12}[g_{i-3,i+3} - 18g_{i-2,i+2} + 63g_{i-1,i+1} - 92g_i], \quad 3 \leq i \leq n-3 \\ \lambda_3 g_i &= g_{i-1} - 2g_i + g_{i+1}, \quad 3 \leq i \leq n-3, \end{aligned}$$

These operators define the relations of eight degree spline S with respect to the higher derivatives $y^{(9)}, \dots, y^{(13)}$.

Lemma 1. *If $y \in C^{14}[a, b]$, then using the above operators we have*

$$\begin{aligned} y_i^{(r)} &= \frac{\lambda_0 S_i^{(r-6)}}{h^6} + \mathcal{O}(h^2), \quad 9 \leq r \leq 13, \quad 3 \leq i \leq n-3, \\ y_i^{(r)} &= \frac{\lambda_1 S_i^{(r-4)}}{h^4} + \mathcal{O}(h^4), \quad 9 \leq r \leq 10, \quad 3 \leq i \leq n-3, \\ y_i^{(11)} &= \frac{\lambda_2 S_i^{(7)}}{h^4} + \mathcal{O}(h^4), \quad 3 \leq i \leq n-3, \\ y_i^{(9)} &= \frac{\lambda_3 S_i^{(7)}}{h^2} + \mathcal{O}(h^6), \quad 1 \leq i \leq n-1. \end{aligned}$$

Proof. The proof is state forward by using Lemma 2.1 and Theorem 2.1. \square

Corollary 1. *Let S be the spline of degree eight which used to interpolate $y \in C^{14}[a, b]$ then for $i = 3(1)n - 3$, the following relations hold*

$$\begin{aligned} y_i^{(7)} &= S_i^{(7)} + \frac{1}{12}\lambda_3 S_i^{(7)} - \frac{1}{240}\lambda_2 S_i^{(7)} + \frac{1}{6048}\lambda_0 S_i^{(7)} + \mathcal{O}(h^8), \\ y_i^{(6)} &= S_i^{(6)} + \frac{1}{240}\lambda_1 S_i^{(6)} - \frac{1}{3024}\lambda_0 S_i^{(6)} + \mathcal{O}(h^8), \\ y_i^{(5)} &= S_i^{(5)} - \frac{1}{720}\lambda_1 S_i^{(5)} + \frac{1}{3024}\lambda_0 S_i^{(5)} + \mathcal{O}(h^8), \\ y_i^{(4)} &= S_i^{(4)} - \frac{1}{6048}\lambda_0 S_i^{(4)} + \mathcal{O}(h^8), \end{aligned}$$

$$\begin{aligned} y_i^{(3)} &= S_i^{(3)} + \frac{1}{30240} \lambda_0 S_i^{(3)} + \mathcal{O}(h^8), \\ y_i^{(2)} &= S_i^{(2)} + \mathcal{O}(h^8), \\ y_i^{(1)} &= S_i^{(1)} + \mathcal{O}(h^8). \end{aligned}$$

Now we need to obtain the similar relations at the boundary and its neighbour points, so that we conclude the following Corollary 2.

Corollary 2. *Let $y \in C^{14}[a, b]$, denoting the index $\sigma_j = j$, $j = 0, 1, 2$ for the grid points, near the left end point and $\sigma_j = n - j$, $j = n - 2, n - 1, n$ for the grid points, near the right end point, then the following approximations to the higher order derivatives of y hold at the boundary and its neighbour points,*

$$\begin{aligned} y_{\sigma_0}^{(r)} &= \lambda_1 \left(\frac{20S_{\sigma_3}^{(r-4)} - 45S_{\sigma_4}^{(r-4)} + 36S_{\sigma_5}^{(r-4)} - 10S_{\sigma_6}^{(r-4)}}{h^4} \right) + \mathcal{O}(h^4), \quad r = 9, 10, \\ y_{\sigma_1}^{(r)} &= \lambda_1 \left(\frac{10S_{\sigma_3}^{(r-4)} - 20S_{\sigma_4}^{(r-4)} + 15S_{\sigma_5}^{(r-4)} - 4S_{\sigma_6}^{(r-4)}}{h^4} \right) + \mathcal{O}(h^4), \quad r = 9, 10, \\ y_{\sigma_2}^{(r)} &= \lambda_1 \left(\frac{4S_{\sigma_3}^{(r-4)} - 6S_{\sigma_4}^{(r-4)} + 4S_{\sigma_5}^{(r-4)} - S_{\sigma_6}^{(r-4)}}{h^4} \right) + \mathcal{O}(h^4), \quad r = 9, 10, \\ y_{\sigma_0}^{(9)} &= \lambda_3 \left(\frac{6S_{\sigma_1}^{(7)} - 15S_{\sigma_2}^{(7)} + 20S_{\sigma_3}^{(7)} - 15S_{\sigma_4}^{(7)} + 6S_{\sigma_5}^{(7)} - S_{\sigma_6}^{(7)}}{h^2} \right) + \mathcal{O}(h^6), \\ y_{\sigma_0}^{(11)} &= \lambda_2 \left(\frac{20S_{\sigma_3}^{(7)} - 45S_{\sigma_4}^{(7)} + 36S_{\sigma_5}^{(7)} - 10S_{\sigma_6}^{(7)}}{h^4} \right) + \mathcal{O}(h^4), \\ y_{\sigma_1}^{(11)} &= \lambda_2 \left(\frac{10S_{\sigma_3}^{(7)} - 20S_{\sigma_4}^{(7)} + 15S_{\sigma_5}^{(7)} - 4S_{\sigma_6}^{(7)}}{h^4} \right) + \mathcal{O}(h^4), \\ y_{\sigma_2}^{(11)} &= \lambda_2 \left(\frac{4S_{\sigma_3}^{(7)} - 6S_{\sigma_4}^{(7)} + 4S_{\sigma_5}^{(7)} - S_{\sigma_6}^{(7)}}{h^4} \right) + \mathcal{O}(h^4), \\ y_{\sigma_0}^{(r)} &= \lambda_0 \left(\frac{4S_{\sigma_3}^{(r-6)} - 3S_{\sigma_4}^{(r-6)}}{h^6} \right) + \mathcal{O}(h^2), \quad r = 9, 10, 11, 12, 13, \\ y_{\sigma_1}^{(r)} &= \lambda_0 \left(\frac{3S_{\sigma_3}^{(r-6)} - 2S_{\sigma_4}^{(r-6)}}{h^6} \right) + \mathcal{O}(h^2), \quad r = 9, 10, 11, 12, 13, \\ y_{\sigma_2}^{(r)} &= \lambda_0 \left(\frac{2S_{\sigma_3}^{(r-6)} - S_{\sigma_4}^{(r-6)}}{h^6} \right) + \mathcal{O}(h^2), \quad r = 9, 10, 11, 12, 13. \end{aligned}$$

3 Description of the method

For solution of system of boundary value problems (1)–(2) by using $S(x) \in Sp_8(\Delta)$ and to achieve an $\mathcal{O}(h^8)$ optimal order method. We approximate

$y', \dots, y^{(7)}$ by their spline relations, which prescribe in Theorem 1, Lemma 1 and Corollaries 1 and 2. Finally this approach lead to the following nonlinear system:

$$\begin{aligned}
& S_{\sigma_0}^{(7)} + \frac{\lambda_3}{12}(6S_{\sigma_1}^{(7)} - 15S_{\sigma_2}^{(7)} + 20S_{\sigma_3}^{(7)} - 15S_{\sigma_4}^{(7)} + 6S_{\sigma_5}^{(7)} - S_{\sigma_6}^{(7)}) \\
& \quad - \frac{\lambda_2}{240}(20S_{\sigma_3}^{(7)} - 45S_{\sigma_4}^{(7)} + 36S_{\sigma_5}^{(7)} - 10S_{\sigma_6}^{(7)}) + \frac{\lambda_0}{6048}(4S_{\sigma_3}^{(7)} - 3S_{\sigma_4}^{(7)}) \\
& = f(x_{\sigma_0}, S_{\sigma_0}, S'_{\sigma_0}, S''_{\sigma_0}, S_{\sigma_0}^{(3)}) + \frac{\lambda_0}{30240}(4S_{\sigma_3}^{(3)} - 3S_{\sigma_4}^{(3)}), \\
& \quad S_{\sigma_0}^{(4)} - \frac{\lambda_0}{6048}(4S_{\sigma_3}^{(4)} - 3S_{\sigma_4}^{(4)}), \\
& \quad S_{\sigma_0}^{(5)} - \frac{\lambda_1}{720}(20S_{\sigma_3}^{(5)} - 45S_{\sigma_4}^{(5)} + 36S_{\sigma_5}^{(5)} - 10S_{\sigma_6}^{(5)}) + \frac{\lambda_0}{3024}(4S_{\sigma_3}^{(5)} - 3S_{\sigma_4}^{(5)}), \\
& \quad S_{\sigma_0}^{(6)} + \frac{\lambda_1}{240}(20S_{\sigma_3}^{(6)} - 45S_{\sigma_4}^{(6)} + 36S_{\sigma_5}^{(6)} - 10S_{\sigma_6}^{(6)}) - \frac{\lambda_0}{3024}(4S_{\sigma_3}^{(6)} - 3S_{\sigma_4}^{(6)}) \\
& \quad + \mathcal{O}(h^8), \quad i = 0, n, \tag{13}
\end{aligned}$$

$$\begin{aligned}
& S_{\sigma_1}^{(7)} + \frac{\lambda_3}{12}S_{\sigma_1}^{(7)} - \frac{\lambda_2}{240}(10S_{\sigma_3}^{(7)} - 20S_{\sigma_4}^{(7)} + 15S_{\sigma_5}^{(7)} - 4S_{\sigma_6}^{(7)}) \\
& \quad + \frac{\lambda_0}{6048}(3S_{\sigma_3}^{(7)} - 2S_{\sigma_4}^{(7)}) = f(x_{\sigma_1}, S_{\sigma_1}, S'_{\sigma_1}, S''_{\sigma_1}), \\
& \quad S_{\sigma_1}^{(3)} + \frac{\lambda_0}{30240}(3S_{\sigma_3}^{(3)} - 2S_{\sigma_4}^{(3)}), S_{\sigma_1}^{(4)} - \frac{\lambda_0}{6048}(3S_{\sigma_3}^{(4)} - 2S_{\sigma_4}^{(4)}), \\
& \quad S_{\sigma_1}^{(5)} - \frac{\lambda_1}{720}(10S_{\sigma_3}^{(5)} - 20S_{\sigma_4}^{(5)} + 15S_{\sigma_5}^{(5)} - 4S_{\sigma_6}^{(5)}) + \frac{\lambda_0}{3024}(3S_{\sigma_3}^{(5)} - 2S_{\sigma_4}^{(5)}), \\
& \quad S_{\sigma_1}^{(6)} + \frac{\lambda_1}{240}(10S_{\sigma_3}^{(6)} - 20S_{\sigma_4}^{(6)} + 15S_{\sigma_5}^{(6)} - 4S_{\sigma_6}^{(6)}) - \frac{\lambda_0}{3024}(3S_{\sigma_3}^{(6)} - 2S_{\sigma_4}^{(6)}) \\
& \quad + \mathcal{O}(h^8), \quad i = 1, n - 1, \tag{14}
\end{aligned}$$

$$\begin{aligned}
& S_{\sigma_2}^{(7)} + \frac{\lambda_3}{12}S_{\sigma_2}^{(7)} - \frac{\lambda_2}{240}(4S_{\sigma_3}^{(7)} - 6S_{\sigma_4}^{(7)} + 4S_{\sigma_5}^{(7)} - S_{\sigma_6}^{(7)}) + \frac{\lambda_0}{6048}(2S_{\sigma_3}^{(7)} - S_{\sigma_4}^{(7)}) \\
& = f(x_{\sigma_2}, S_{\sigma_2}, S'_{\sigma_2}, S''_{\sigma_2}, S_{\sigma_2}^{(3)}) + \frac{\lambda_0}{30240}(2S_{\sigma_3}^{(3)} - S_{\sigma_4}^{(3)}), \\
& \quad S_{\sigma_2}^{(4)} - \frac{\lambda_0}{6048}(2S_{\sigma_3}^{(4)} - S_{\sigma_4}^{(4)}), \\
& \quad S_{\sigma_2}^{(5)} - \frac{\lambda_1}{720}(4S_{\sigma_3}^{(5)} - 6S_{\sigma_4}^{(5)} + 4S_{\sigma_5}^{(5)} - S_{\sigma_6}^{(5)}) + \frac{\lambda_0}{3024}(2S_{\sigma_3}^{(5)} - S_{\sigma_4}^{(5)}), \\
& \quad S_{\sigma_2}^{(6)} + \frac{\lambda_1}{240}(4S_{\sigma_3}^{(6)} - 6S_{\sigma_4}^{(6)} + 4S_{\sigma_5}^{(6)} - S_{\sigma_6}^{(6)}) - \frac{\lambda_0}{3024}(2S_{\sigma_3}^{(6)} - S_{\sigma_4}^{(6)}) \\
& \quad + \mathcal{O}(h^8), \quad i = 2, n - 2, \tag{15}
\end{aligned}$$

$$\begin{aligned}
S_i^{(7)} + \frac{\lambda_3}{12} S_i^{(7)} - \frac{\lambda_2}{240} S_i^{(7)} + \frac{\lambda_0}{6048} S_i^{(7)} &= f(x_i, S_i, S_i', S_i'', S_i^{(3)}) + \frac{\lambda_0}{30240} S_i^{(3)}, \\
S_i^{(4)} - \frac{\lambda_0}{6048} S_i^{(4)}, S_i^{(5)} - \frac{\lambda_1}{720} S_i^{(5)} + \frac{\lambda_0}{3024} S_i^{(5)}, \\
S_i^{(6)} + \frac{\lambda_1}{240} S_i^{(6)} - \frac{\lambda_0}{3024} S_i^{(6)} &+ \mathcal{O}(h^8), \quad 3 \leq i \leq n-3,
\end{aligned} \tag{16}$$

associated with the boundary formulas,

$$\begin{aligned}
BS \equiv & \alpha_{i,0} S_0 + \alpha_{i,1} S_0' + \alpha_{i,2} S_0'' + \alpha_{i,3} (S_0^{(3)} + \frac{\lambda_0}{30240} (4S_3^{(3)} - 3S_4^{(3)})) \\
& + \alpha_{i,4} (S_0^{(4)} - \frac{\lambda_0}{6048} (4S_3^{(4)} - 3S_4^{(4)})) \\
& + \alpha_{i,5} (S_0^{(5)} - \frac{\lambda_1}{720} (20S_{\sigma_3}^{(5)} - 45S_{\sigma_4}^{(5)} + 36S_{\sigma_5}^{(5)} - 10S_{\sigma_6}^{(5)})) \\
& + \frac{\lambda_0}{3024} (4S_3^{(5)} - 3S_4^{(5)}) \\
& + \alpha_{i,6} (S_0^{(6)} + \frac{\lambda_1}{240} (20S_{\sigma_3}^{(6)} - 45S_{\sigma_4}^{(6)} + 36S_{\sigma_5}^{(6)} - 10S_{\sigma_6}^{(6)})) \\
& - \frac{\lambda_0}{3024} (4S_3^{(6)} - 3S_4^{(6)}) + \beta_{i,0} S_n + \beta_{i,1} S_n' + \beta_{i,2} S_n'' \\
& + \beta_{i,3} (S_n^{(3)} + \frac{\lambda_0}{30240} (4S_{n-3}^{(3)} - 3S_{n-4}^{(3)})) \\
& + \beta_{i,4} (S_n^{(4)} - \frac{\lambda_0}{6048} (4S_{n-3}^{(4)} - 3S_{n-4}^{(4)})) \\
& + \beta_{i,5} (S_n^{(5)} - \frac{\lambda_1}{720} (20S_{n-3}^{(5)} - 45S_{n-4}^{(5)} + 36S_{n-5}^{(5)} - 10S_{n-6}^{(5)})) \\
& + \frac{\lambda_0}{3024} (4S_{n-3}^{(5)} - 3S_{n-4}^{(5)}) \\
& + \beta_{i,6} (S_n^{(6)} + \frac{\lambda_1}{240} (20S_{n-3}^{(6)} - 45S_{n-4}^{(6)} + 36S_{n-5}^{(6)} - 10S_{n-6}^{(6)})) \\
& - \frac{\lambda_0}{3024} (4S_{n-3}^{(6)} - 3S_{n-4}^{(6)}) = \eta_i, \quad i = 0, 1, \dots, 6.
\end{aligned} \tag{17}$$

Let L' be the approximation of L defined as follows,

$$\begin{aligned}
L'g_i \equiv & g_i^{(7)} + \frac{1}{12} \lambda_3 g_i^{(7)} \\
& - \frac{1}{240} \lambda_2 g_i^{(7)} + \frac{1}{6048} \lambda_0 g_i^{(7)} - f(x_i, g_i, g_i', g_i'', g_i^{(3)}) + \frac{1}{30240} \lambda_0 g_i^{(3)}, \\
& g_i^{(4)} - \frac{1}{6048} \lambda_0 g_i^{(4)}, g_i^{(5)} - \frac{1}{720} \lambda_1 g_i^{(5)} + \frac{1}{3024} \lambda_0 g_i^{(5)}, \\
& g_i^{(6)} + \frac{1}{240} \lambda_1 g_i^{(6)} - \frac{1}{3024} \lambda_0 g_i^{(6)},
\end{aligned}$$

and let B' be the approximation of B defined in (17), and $S(x)$ be the spline of degree eight which is the solution of the system (1) – (2), then the following relations hold,

$$\begin{cases} L'S_i = \mathcal{O}(h^8), & 0 \leq i \leq n, \\ B'S = \mathcal{O}(h^8). \end{cases} \quad (18)$$

For the convergence analysis first we need to recall and prove the following Lemmas.

Lemma 2. *If $p = \{p_{ij}\}$ is an $m \times m$ matrix and $p_{ii} \geq \sum_{j=1, i \neq j}^m |p_{ij}| + \epsilon$, for $i = 1, 2, \dots, m$, where $\epsilon > 0$, then we have $\|p^{-1}\|_\infty \leq \epsilon^{-1}$.*

Proof. See Lemma 4 in [13]. \square

Lemma 3. *If the coefficients matrix of $S_i^{(7)}$ in the equation $L'S_i = \mathcal{O}(h^8)$, $i = 0, 1, \dots, n$, is denoted by Q_7 , then Q_7 is nonsingular and $\|Q_7^{-1}\|_\infty$ is bounded.*

Proof. Using relations (13) – (16) we can obtain the $(n+1) \times (n+1)$ coefficients matrix Q_7 as

$$Q_7 = \frac{1}{60480} \times$$

$$\begin{pmatrix} 91180 & -144855 & 327246 & -466043 & \dots & 4482248 & -291687 & 124918 & -32853 \\ 5280 & 46000 & 26715 & -52434 & \dots & 74077 & -64992 & 35493 & -11402 \\ 104 & 3272 & 58404 & -12709 & \dots & 23054 & -18771 & 9656 & -2971 \\ 31 & -438 & 6513 & 48268 & \dots & 6513 & -438 & 31 & 0 \\ 0 & 31 & -438 & 6513 & \dots & 48268 & 6513 & -438 & 31 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 31 & -438 & \dots & 6513 & 48268 & 6513 & -438 \\ 0 & 0 & 0 & 31 & \dots & -438 & 6513 & 48268 & 6513 \\ -21 & 4625 & -2971 & 9656 & \dots & -18771 & 23054 & -12709 & 58404 \\ -84 & 1827 & -11402 & 35493 & \dots & -64992 & 74077 & -52434 & 26715 \\ -210 & 4536 & -32853 & 124918 & \dots & -291687 & 4482248 & -466043 & 327246 \end{pmatrix},$$

Let E_i , be the i -th row of Q_7 . Then by using the following elementary row operations, this matrix can be converted to a strictly diagonally dominant:

$$\begin{aligned} E_2 + \frac{1}{2}E_3 - \frac{1}{4}E_4 + \frac{1}{4}E_5 &\rightarrow E_2, \\ E_1 - \frac{1}{2}E_2 + E_3 - 1.4E_4 + 1.35E_5 - 0.833E_6 + 0.333E_7 &\rightarrow E_1, \\ 0.1E_0 + 0.314E_1 - 0.847E_2 + 1.61E_3 - 1.56E_4 + 1.172E_5 \\ &\quad - 0.4859E_6 + 0.1666E_7 + 0.0407E_8 \rightarrow E_0, \\ E_{n-2} + \frac{1}{2}E_{n-3} - \frac{1}{4}E_{n-4} + \frac{1}{4}E_{n-5} &\rightarrow E_{n-2}, \end{aligned}$$

$$\begin{aligned}
& E_{n-1} - \frac{1}{2}E_{n-2} + E_{n-3} - 1.4E_{n-4} + 1.35E_{n-5} - 0.833E_{n-6} \\
& \quad + 0.333E_{n-7} \rightarrow E_{n-1}, \\
& 0.1E_n + 0.314E_{n-1} - 0.847E_{n-2} + 1.61E_{n-3} - 1.56E_{n-4} + 1.172E_{n-5} \\
& \quad - 0.4859E_{n-6} + 0.1666E_{n-7} + 0.0407E_{n-8} \rightarrow E_n,
\end{aligned}$$

Hence, the matrix Q_7 is strictly diagonally dominant and positive definite. Therefore, using Lemma 2 we can conclude that $\|Q_7^{-1}\|_\infty$ is finite. \square

4 Convergence analysis

We prove the convergence of the presented method via Green's function scheme. If we assume that the boundary value condition $y^{(7)} = 0$ subjected to homogeneous boundary conditions $By = 0$, has a unique solution, it implies that there is a Green's function $G(x, t)$ for this problem [19]. Let $y^{(7)} = \phi$ and $\hat{S}^{(7)} = \psi$, be the exact and the spline solutions of the problem (1) which satisfy the boundary conditions (2). Then $y(x)$ and $\hat{S}(x)$ and its first sixth derivatives can be obtained as follows:

$$y^{(i)}(x) = \int_a^b \frac{\partial^i G(x, t)}{\partial x^i} \phi(t) dt, \quad \hat{S}^{(i)}(x) = \int_a^b \frac{\partial^i G(x, t)}{\partial x^i} \psi(t) dt,$$

for $i = 0, 1, \dots, 6$. We define the operators $F_n, \mathbf{M}_n, \mathbf{k}$ and \mathbf{R} as:

$$F_n : C[a, b] \rightarrow R^{n+1}, \quad F_n g = [g(x_0), \dots, g(x_n)]^T,$$

$$\mathbf{M}_n : R^{n+1} \rightarrow C[a, b], \quad \text{via piecewise linear interpolation at } \{x_i\}_0^n,$$

$$\mathbf{k} : C[a, b] \rightarrow C[a, b], \quad \mathbf{k}g = f(x, G_{p,0}(x), G_{p,1}(x), \dots, G_{p,6}(x)),$$

$$\mathbf{R} : C[a, b] \rightarrow C[a, b], \quad \mathbf{R}g = f(x, Q_0 F_n G_{p,0}(x), \dots, Q_6 F_n G_{p,6}(x)),$$

where $g \in C[a, b]$, $G_{p,i}(x) = \int_a^b \frac{\partial^i G_p(x, t)}{\partial x^i} g(t) dt$, $i = 0, 1, \dots, 6$ and

$$Q_i = \begin{cases} I_{(n+1) \times (n+1)}, & 0 \leq i \leq 2, \\ \text{The coefficients matrix of } S^{(i)} \text{ in Eq. (18)}, & 3 \leq i \leq 7. \end{cases}$$

With the introduced notations, we can rewrite Eqs. (1) and (13)-(16) respectively as:

$$y^{(7)} - f(x, y(x), y'(x), \dots, y^{(6)}(x)) = \phi - \mathbf{k}\phi = (I - \mathbf{k})\phi = 0, \quad (21)$$

$$Q_7 F_n \hat{S}^{(7)} - f(x, Q_0 F_n \hat{S}, Q_1 F_n \hat{S}', \dots, Q_6 F_n \hat{S}^{(6)}) = Q_7 F_n \hat{S}^{(7)} - F_n \mathbf{R} \psi = 0.$$

Since Q_7 is nonsingular and $\hat{S}^{(7)}(x)$ is a linear polynomial, therefore we have the following relations:

$$F_n \hat{S}^{(7)} - Q_7^{-1} F_n \mathbf{R} \psi = 0 \Rightarrow \mathbf{M}_n F_n \hat{S}^{(7)} - \mathbf{M}_n Q_7^{-1} F_n \mathbf{R} \psi = 0,$$

$$\hat{S}^{(7)} - \mathbf{M}_n Q_7^{-1} F_n \mathbf{R} \psi = (I - \mathbf{M}_n Q_7^{-1} F_n \mathbf{R}) \psi = (I - p_n \mathbf{R}) \psi = 0, \quad (22)$$

where $p_n = \mathbf{M}_n Q_7^{-1} F_n$. Notice that p_n is an operator from $C[a, b]$ into the continuous piecewise linear functions with grid points x_i .

Lemma 4. *Let $\{\Delta\}$ be a sequence of partitions of the interval $[a, b]$. Then the sequence of operators $p_n = \mathbf{M}_n Q_7^{-1} F_n$ converges to the identity operator as h approaches zero.*

Proof. We want to show that $|p_n g - g| \rightarrow 0$ for each $g \in C[a, b]$. To do so, we have

$$\begin{aligned} \|p_n g - g\| &\leq \|\mathbf{M}_n Q_7^{-1} F_n g - \mathbf{M}_n F_n g\| \\ &\leq \|\mathbf{M}_n\| \|Q_7^{-1}\| \|F_n g - Q_7 F_n g\| \\ &\leq C^* \|F_n g - Q_7 F_n g\| \\ &\leq C^* \omega(g, 10h), \end{aligned}$$

where C^* is a finite constant and $\omega(g, \epsilon) = \sup\{|g(x + \epsilon') - g(x)| : x, x + \epsilon' \in [a, b], |\epsilon'| \leq \epsilon\}$. When $h \rightarrow 0$ we have, $\omega(g, 10h) \rightarrow 0$. \square

Lemma 5. *Let $g \in C[a, b]$, then $p_n \mathbf{R}$ converges to \mathbf{k} .*

Proof. By using the definitions of \mathbf{k} and \mathbf{R} we obtain

$$\begin{aligned} \|p_n \mathbf{R} g - \mathbf{k} g\| &= \|\mathbf{M}_n Q_7^{-1} F_n \mathbf{R} g - \mathbf{k} g\| \\ &\leq \|\mathbf{M}_n Q_7^{-1} F_n \mathbf{R} g - \mathbf{M}_n F_n \mathbf{k} g\| + \|\mathbf{M}_n F_n \mathbf{k} g - \mathbf{k} g\| \\ &\leq \|\mathbf{M}_n Q_7^{-1}\| \|F_n \mathbf{R} g - Q_7 F_n \mathbf{K} g\| + \mathcal{O}(h^2), \end{aligned}$$

and since $\|\mathbf{M}_n\|$ and $\|Q_7^{-1}\|$ are bounded, we have

$$\|p_n \mathbf{R} g - \mathbf{k} g\| \leq \hat{C} \|F_n \mathbf{R} g - Q_7 F_n \mathbf{K} g\| \leq \hat{C} \omega(g, 10\delta),$$

with

$$\delta = \max\{10h, \omega(G_{p,0}(x), 17h), \omega(G_{p,1}(x), 17h), \dots, \omega(G_{p,6}(x), 17h)\}. \quad (23)$$

$\omega(G_{p,j}(x), 17h)$, $0 \leq j \leq 6$ convergence to zero as h approaches zero for continuous functions $G_{p,j}(x)$, $0 \leq j \leq 6$, so that by using Eq (23), $\delta \rightarrow 0$ and $\omega(g, 10\delta)$ convergence to zero. \square

Now we present the main convergence theorem.

Theorem 2. *The error bounds for collocation approximation $\hat{S}(x) \in Sp_8(\Delta)$ satisfies,*

$$\begin{aligned} \|(y - \hat{S})^{(j)}\| &= \mathcal{O}(h^{8-j}), \quad j = 0, 1, \dots, 7, \\ |(y - \hat{S})_i^{(j)}| &= \mathcal{O}(h^8), \quad j = 0, 1, 2, \\ |(y - \hat{S})_i^{(j)}| &= \mathcal{O}(h^6), \quad j = 3, 4, \\ |(y - \hat{S})_i^{(j)}| &= \mathcal{O}(h^4), \quad j = 5, 6, \\ |(y - \hat{S})_i^{(j)}| &= \mathcal{O}(h^2), \quad j = 7. \end{aligned}$$

Proof. We consider the problem: $S^{(7)} = \nu$, $BS = \mathcal{O}(h^8)$. Let $\{\Delta\}$ be a sequence of partitions of the $[a, b]$ and the problem $y^{(7)} = 0$, $By = 0$ has a unique solution. So there exists a polynomial $\xi(x)$ of order 6 as follows

$$B\xi = BS = \mathcal{O}(h^8), \quad \|\xi^{(k)}\| = \mathcal{O}(h^8), \quad k = 0, 1, \dots, 6. \quad (24)$$

From solvability of $(S - \xi)^{(7)} = \nu$, $B(S - \xi) = 0$ we obtain

$$(I - \mathbf{M}_n Q_7^{-1} F_n \mathbf{R})(S^{(7)} - \xi^{(7)}) = \mathbf{M}_n Q_7^{-1} (Q_7 F_n - F_n \mathbf{R})(S - \xi)^{(7)}.$$

Using (18) and the boundedness of $\|\mathbf{M}_n\|$ and $\|Q_7^{-1}\|$, we have

$$(I - \mathbf{M}_n Q_7^{-1} F_n \mathbf{R})(S^{(7)} - \xi^{(7)}) = \mathbf{M}_n Q_7^{-1} (\mathcal{O}(h^8)) = \mathcal{O}(h^8). \quad (25)$$

Subtracting (22) and (25), we obtain,

$$(I - \mathbf{M}_n Q_7^{-1} F_n \mathbf{R})(S^{(7)} - \xi^{(7)} - \hat{S}^{(7)}) = \mathcal{O}(h^8),$$

and we have

$$(S^{(7)} - \xi^{(7)} - \hat{S}^{(7)}) = p_n \mathbf{R}(S^{(7)} - \xi^{(7)} - \hat{S}^{(7)}) + \mathcal{O}(h^8). \quad (26)$$

The operator \mathbf{R} is continuously differentiable. So Eq. (26) has an integral equation form as following

$$\begin{aligned} (S^{(7)} - \xi^{(7)} - \hat{S}^{(7)}) &= p_n \left(\int_0^1 (\mathbf{R}'[\hat{S}^{(7)} + t(S^{(7)} - \xi^{(7)} - \hat{S}^{(7)})] dt) \right) \\ &\quad \times (S^{(7)} - \xi^{(7)} - \hat{S}^{(7)}) + \mathcal{O}(h^8), \end{aligned} \quad (27)$$

where $\{\tau_n\} = p_n \left(\int_0^1 (\mathbf{R}'[\psi + t(S^{(7)} - \xi^{(7)} - \hat{S}^{(7)})] dt) \right)$, is a sequence of linear operators converging to $\mathbf{R}'(y^{(7)})$. So we have

$$(S^{(7)} - \xi^{(7)} - \hat{S}^{(7)}) = \tau_n (S^{(7)} - \xi^{(7)} - \hat{S}^{(7)}) + \mathcal{O}(h^8).$$

Since $(I - \tau_n)^{-1}$ exists and its norm is bounded, we obtain

$$\|(S - \xi - \hat{S})^{(7)}\|_\infty = \mathcal{O}(h^8). \quad (28)$$

According to the hypotheses of the problem, $(S - \xi - \hat{S})^{(7)} = r$, $B(S - \xi - \hat{S}) = 0$, has unique solution. So we can write $(S - \xi - \hat{S})^{(i)}$ in the following form

$$(S - \xi - \hat{S})^{(i)} = \int \frac{\partial^i G(x, t)}{\partial x^i} (S^{(7)} - \xi^{(7)} - \hat{S}^{(7)})(t) dt, \quad i = 0, 1, \dots, 6, \quad (29)$$

which implies that

$$\|(S - \xi - \hat{S})^{(i)}\|_\infty = \mathcal{O}(h^8), \quad i = 0, 1, \dots, 6. \quad (30)$$

Using the triangular inequality we obtain

$$\|(y - S)^{(i)}\| \leq \|(y - \hat{S})^{(i)}\| + \|(S - \hat{S})^{(i)}\| + \|\xi^{(i)}\|, \quad i = 0, 1, \dots, 6,$$

by using equations (18) and (24) and Theorem 1, we can obtain the results of Theorem 2. This completes of proof. \square

5 Numerical experiments

We present the results from numerical experiments to demonstrate the performance of the presented method and verify the results of the analysis. The obtained results has been compared with the references [21, 23, 3] and the results tabulated in Tables 1–6, these results verify the accurate nature of our purposed method in applications. The numerical computations have done by the software Mathematica 10.

Example 1. The following linear seventh order boundary value problem is considered:

$$y^{(7)}(x) = xy(x) + e^x(x^2 - 2x - 6), \quad 0 \leq x \leq 1,$$

subjected to the boundary conditions

$$\begin{aligned} y(0) = y(1) = 1, \quad y'(0) = 0, \quad y'(1) = -e, \\ y''(0) = -1, \quad y''(1) = -2e, \quad y^{(3)}(0) = -2. \end{aligned}$$

The exact solution of the problem is $y(x) = (1 - x)e^x$. This example has been solved by our method with $h = \frac{1}{10}$, the maximum absolute errors in the certain points are tabulated in Table 1 and compared with [21, 23],

which shows that our method is accurate. Also the example has been solved with $h = \frac{1}{9}, \frac{1}{18}, \frac{1}{36}, \frac{1}{72}, \frac{1}{144}$ and the maximum absolute errors in the solutions are tabulated in Table 2. In this table $E_i = \|y^{(i)} - \hat{S}^{(i)}\|_{\infty}$, $0 \leq i \leq 6$ and O_i is the order of convergence of i -th derivatives of y . This table also verified that our approach are applicable and accurate.

Table 1: The maximum absolute errors in the solution of Example 1.

x	our method	method in [23]	method in [21]
0.1	1.03(-15)	3.42(-13)	4.66(-13)
0.2	2.08(-15)	6.25(-14)	5.71(-12)
0.3	5.71(-14)	1.42(-13)	2.13(-11)
0.4	9.52(-14)	8.84(-14)	4.69(-11)
0.5	8.82(-14)	6.43(-14)	7.43(-11)
0.6	5.05(-13)	1.52(-12)	8.92(-11)
0.7	1.91(-13)	1.48(-12)	7.98(-11)
0.8	1.82(-13)	4.94(-12)	4.67(-11)
0.9	1.05(-13)	5.38(-12)	1.09(-11)

Table 2: The maximum absolute errors in the solution of Example 1 with various values of h .

h	$\frac{1}{9}$	$\frac{1}{18}$	$\frac{1}{36}$	$\frac{1}{72}$	$\frac{1}{144}$
E_0, O_0	1.5(-13), -	5.3(-16), 8.1	1.9(-18), 8.1	7.6(-21), 7.9	3.0(-23), 7.9
E_1, O_1	5.6(-13), -	2.0(-15), 8.1	7.8(-18), 8	2.9(-20), 8.1	1.1(-22), 8
E_2, O_2	4.2(-12), -	1.6(-14), 8	5.9(-17), 8.1	2.3(-19), 8	8.9(-22), 8
E_3, O_3	8.9(-10), -	1.5(-11), 5.9	2.3(-13), 6	3.7(-15), 6	5.7(-17), 6
E_4, O_4	5.0(-9), -	8.1(-11), 5.9	1.3(-12), 5.9	2.0(-14), 6	3.2(-16), 6
E_5, O_5	3.1(-6), -	1.9(-7), 4	1.3(-8), 3.9	7.9(-10), 4	4.9(-11), 4
E_6, O_6	1.0(-5), -	6.6(-7), 3.9	4.2(-8), 4	2.7(-9), 4	1.8(-10), 4

Example 2. Consider the following nonlinear seventh order boundary value problem,

$$y^{(7)}(x) = y^2(x)e^x, \quad 0 \leq x \leq 1,$$

subjected to the boundary conditions

$$y(0) = y'(0) = y''(0) = y^{(3)}(0) = 1, \quad y(1) = y'(1) = y''(1) = e.$$

Table 3: The maximum absolute errors in the solution of Example 2.

h	our method	method in [3]	method in [21]
$\frac{1}{10}$	6.22(-15)	6.48(-11)	3.02(-14)
$\frac{1}{30}$	1.42(-18)	3.31(-14)	--
$\frac{1}{50}$	1.01(-19)	2.78(-15)	--

The exact solution of this problem is $y(x) = e^x$. First of all we solve this problem for various values of $h = \frac{1}{10}, \frac{1}{30}, \frac{1}{50}$ and compare with the results in [21, 3]. Our results are shown in Table 3. Then we obtain E_i and O_i for various values of h . The results are tabulated in Table 4. This table shows that the orders of convergence in applications agree with those we obtained theoretically.

Table 4: The maximum absolute errors in the solution of Example 2 with various values of h .

h	$\frac{1}{9}$	$\frac{1}{18}$	$\frac{1}{36}$	$\frac{1}{72}$	$\frac{1}{144}$
E_0, O_0	1.5(-14), -	5.3(-17), 8.1	2.0(-19), 8	7.7(-22), 8	2.9(-24), 8.1
E_1, O_1	5.9(-14), -	2.2(-16), 8.1	8.3(-19), 8.1	3.2(-21), 8	1.2(-23), 8.1
E_2, O_2	4.3(-13), -	1.6(-15), 8.1	6.2(-18), 8	2.4(-20), 8	9.3(-23), 8
E_3, O_3	1.0(-10), -	1.7(-12), 5.9	2.7(-14), 6	4.2(-16), 6	6.6(-18), 6
E_4, O_4	5.2(-10), -	8.4(-12), 6	1.3(-13), 6	2.1(-15), 6	3.3(-17), 6
E_5, O_5	3.6(-7), -	2.3(-8), 4	1.5(-9), 3.9	9.2(-11), 4	5.8(-12), 4
E_6, O_6	1.1(-6), -	6.9(-8), 4	4.4(-9), 4	2.8(-10), 4	1.7(-11), 4

Example 3. Consider the following nonlinear seventh order boundary value problem,

$$y^{(7)}(x) + y^{(4)}(x) - y(x)e^{y(x)} = e^x((-4(-3+x) + e^{(-e^x(x-1)\cos x)}(x-1))\cos x - 8(5+x)\sin x), \quad 0 \leq x \leq 1,$$

subjected to the boundary conditions

$$y(0) = 1, \quad y'(0) = y(1) = 0, \quad y'(1) = -e \cos 1, \\ y''(0) = y^{(3)}(0) = -2, \quad y''(1) = -2e \cos 1 + 2e \sin 1.$$

The exact solution of the problem is given by $y(x) = e^x(1-x)\cos x$. First we solve this problem with $h = \frac{1}{30}$ and compared the errors in those special

points given in [3]. These results are tabulated in Table 5, the results in this table verified that our method is more accurate. Then we obtain E_i and O_i for various values of h . The results are tabulated in Table 6.

Table 5: The maximum absolute errors in the solution of Example 3.

x	our method	method in [3]
0.125	1.15(-12)	4.74(-10)
0.250	1.35(-10)	5.20(-9)
0.375	2.89(-9)	1.53(-8)
0.500	5.42(-9)	2.45(-8)
0.625	4.96(-9)	2.53(-8)
0.750	2.69(-9)	1.56(-8)
0.875	1.94(-10)	3.29(-9)

Table 6: The maximum absolute errors in the solution of Example 3 with various values of h .

h	$\frac{1}{9}$	$\frac{1}{18}$	$\frac{1}{36}$	$\frac{1}{72}$	$\frac{1}{144}$
E_0, O_0	3.8(-13), -	1.1(-15), 8.4	6.3(-18), 7.4	2.9(-20), 7.8	1.2(-22), 7.9
E_1, O_1	4.6(-12), -	1.6(-14), 8.1	6.3(-17), 8	2.5(-19), 8	9.8(-22), 8
E_2, O_2	1.7(-11), -	5.4(-14), 8.3	2.4(-16), 7.8	9.9(-19), 7.9	4.1(-21), 7.9
E_3, O_3	1.2(-8), -	1.9(-10), 6	2.9(-12), 6	4.6(-14), 6	7.2(-16), 6
E_4, O_4	3.9(-8), -	5.8(-10), 6.1	9.1(-12), 6	1.4(-13), 6	2.2(-15), 6
E_5, O_5	4.1(-5), -	2.6(-6), 4	1.6(-7), 4	1.1(-8), 3.9	6.3(-10), 4.1
E_6, O_6	7.7(-5), -	4.8(-6), 4	2.9(-7), 4	1.8(-8), 4	1.2(-9), 3.9

6 Conclusion

We developed a numerical method to solve the general nonlinear seventh order boundary value problems by using eighth degree B-spline approximation. The numerical illustration shown the proposed method has the $\mathcal{O}(h^8)$ order of accuracy, so we can conclude that our method has highly accurate and efficient in comparison with the other existing methods. Our results obtained by the optimal $\mathcal{O}(h^8)$ method are in good agreement with the proposed numerical algorithm.

Acknowledgements

The authors would like to thank the anonymous referee for the valuable comments.

References

- [1] E.L. Ablasiny and W.D. Hoskins, *Cubic spline solutions to two-point boundary value problems*, Comput. J. **12** (1969) 151-153.
- [2] J.H. Ahlberg, E.N. Nilson and J.L. Walsh, *The Theory of Splines and Their Applications*, Academic Press, New York, 1967.
- [3] G. Akram and H.U. Rehman, *Numerical solution of seventh order boundary value problems using the reproducing kernel space*, Res. J. App. Sci. Eng. and Technol. **7** (2014) 892–896.
- [4] A. Bellen and M. Zennaro, *A collocation method for boundary value problems of differential equations with functional arguments*, Computing **32** (1984) 307-318.
- [5] W.G. Bickley, *Piecewise cubic interpolation and two-point boundary value problems*, Comput. J. **11** (1968) 206-208.
- [6] J.W. Daniel and B.K. Swartz, *Extrapolated collocation for two-point boundary value problems using cubic splines*, J. Inst. Math. Appl. **16** (1975) 161–174.
- [7] D.J. Fyfe, *The use of cubic splines in the solution of two-point boundary value problems*, Comput. J. **12**(1969) 188-192.
- [8] D.J. Fyfe, *The use of cubic splines in the solution of certain fourth order boundary value problems*, Comput. J. **13** (1970) 204-205.
- [9] D.J. Fyfe, *Linear dependence relations connecting equal interval n th degree spline and their derivatives*, J. Inst. Math. Appl. **7** (1971) 398-406.
- [10] E.N. Houstis, C.C. Christara and J.R. Rice, *Quadratic spline collocation methods for two-point boundary value problems*, Int. J. Numer. Methods Eng. **26** (1988) 935-952.
- [11] M. Irodotou-Ellina and E.N. Houstis, *An $\mathcal{O}(h^6)$ quintic spline collocation method for fourth order two-point boundary value problems*, BIT Numerical Mathematics. **28**(1988) 288-301.

- [12] J.B. Kioustelidis, *An error bounds for approximate solutions of two point boundary value problems*, Computing. **42**(1989) 259-270.
- [13] T.R. Lucas, *Error bounds for interpolating cubic spline under various end conditions*, SIAM J. Numer. Anal. **11** (1974) 569-584.
- [14] R.K. Mohanty, *A fourth-order finite difference method for the general one dimensional nonlinear biharmonic problems of first kind*, J. Comput. Appl. Math. **114**(2000) 275–290.
- [15] J. Rashidinia and M. Nabati, *Sinc-Galerkin and Sinc-Collocation methods in the solution of nonlinear two-point boundary value problems*, Comp. Appl. Math. **32** (2013) 315–330.
- [16] J. Rashidinia, M. Ghasemi and R. Jalilian, *An $\mathcal{O}(h^6)$ numerical solution of general nonlinear fifth-order two point boundary value problems*, Numer. Algor. **55** (2011) 403-428.
- [17] J. Rashidinia and M. Ghasemi, *B-spline collocation for solution of two-point boundary value problems*, J. Comput. Appl. Math. **235** (2011) 2325-2342.
- [18] G. Richards and P. Sarma, *Reduced order models for induction motors with two rotor circuits*, IEEE T. Energy Conver. **9** (1994) 673–678.
- [19] R.D. Russell, L.F. Shampine, *A collocation method for boundary value problems*, Numer. Math. **19** (1972) 1-28.
- [20] M. Sakai, *Piecewise cubic interpolation and two-point boundary value problems*, Publ. Res. Inst. Math. Sci. **7** (1971) 345-362.
- [21] S.S. Siddiqi, G. Akram and M. Iftikhar, *Solution of seventh order boundary value problem by differential transformation method*, World Appl. Sci. J. **16** (2012) 1521–1526.
- [22] S.S. Siddiqi, G. Akram and M. Iftikhar, *Solution of Seventh Order Boundary Value Problems by Variational Iteration Technique*, Appl. Math. Sci. **6** (2012) 4663–4672.
- [23] S.S. Siddiqi and M. Iftikhar, *Numerical solution of higher order boundary value problems*, Abstr. Appl. Anal. **2013** (2013), Article ID 427521.
- [24] S.S. Siddiqi and E.H. Twizell, *Spline solutions of linear eighth order boundary value problems*, Comput. Methods. Appl. Mech. Engrg. **131** (1996) 309–325.