

# Legendre-collocation method to solve the second kind Cauchy integral equations

Arezoo Heidari <sup>†</sup>, Saeid Khezerloo <sup>‡\*</sup>, Alireza Gilani <sup>§</sup>, Rahele Noraei <sup>§</sup> Muhammad Arghand <sup>§</sup>

<sup>†</sup> Department of Mathematics, South Tehran Branch, Islamic Azad University, Tehran, Iran  
Email(s): s\_khezerloo@azad.ac.ir

**Abstract.** A numerical solution for the second kind singular integral equations with Cauchy kernel is developed using the collocation method. To achieve this, we approximate the Cauchy integral equation using the collocation method and Legendre orthogonal polynomial expansions. The accuracy of our proposed method is assessed through convergence and error analysis. Finally, several numerical examples are presented to demonstrate the high efficiency of the method.

*Keywords:* Cauchy integral equation, Legendre polynomials, collocation method.  
*AMS Subject Classification 2010:* 30E20, 45E05, 33C45, 65N35.

## 1 Introduction

Cauchy singular integral equations play a significant role in engineering and applied sciences, both in research and practical applications. These equations are commonly used to describe various phenomena, including electromagnetic radiation, unsteady aerodynamic behaviour, electron microscopy, control problems, viscoelasticity, thermoelasticity, and fluid dynamics [6, 7, 10]. Consider an integral equation that includes the unknown function  $f$  under the integral sign of an improper integral in the sense of Cauchy, represented as follows

$$f(\lambda) = g(\lambda) + \oint_{-1}^1 \frac{f(x)}{\lambda - x} dx, \quad -1 < \lambda < 1, \quad (1)$$

In this context,  $g$  is a known function, and the symbol  $\oint$  denotes the Cauchy principal value integral, which is defined as follows

$$\oint_{-1}^1 = \lim_{\varepsilon \rightarrow 0^+} \left( \int_{-1}^{\lambda - \varepsilon} + \int_{\lambda + \varepsilon}^1 \right).$$

In the special case of the problem mentioned above, the corresponding equation

$$\oint_{-1}^1 \frac{f(x)}{\lambda - x} dx = -g(\lambda), \quad -1 < \lambda < 1,$$

\*Corresponding author

Received: 2 July 2024 / Revised: 29 September 2024 / Accepted: 4 November 2024  
DOI: [10.22124/jmm.2024.27832.2451](https://doi.org/10.22124/jmm.2024.27832.2451)

represents the simplest singular integral equation of the first kind. This equation has a bounded solution, which is discussed in various contexts [3].

Now, consider Eq. (1) and assume that  $f \in H_\alpha[-1, 1]$ , meaning that  $f$  is Hölder's continuous on the interval  $[-1, 1]$ , with Hölder index  $\alpha$ , where  $0 < \alpha \leq 1$ . For all points  $l_1, l_2 \in [-1, 1]$ , the following condition holds

$$|f(l_2) - f(l_1)| \leq M |l_2 - l_1|^\alpha,$$

where  $M$  is independent of  $l_1$  and  $l_2$ . General methods for solving this equation have been studied, particularly regarding its existence and uniqueness. According to references [4, 10], the function  $g$  belongs to  $H_\alpha[-1, 1]$  on  $[-1, 1]$  and we investigate solutions  $f$  that are Hölder continuous on any closed interval within the interior of  $(-1, 1)$  and are integrable at the endpoints  $-1$  and  $+1$ .

In this work, the Collocation method is considered for solving the Cauchy type singular integral equations using Legendre polynomials. Let  $\{p_k(x)\}_{k=0}^\infty$  denote the sequence of Legendre polynomials of degree  $k$  defined on the interval  $[-1, 1]$ . These polynomials satisfy the following recurrence relation [14]

$$\begin{aligned} p_0(x) &= 1, \quad p_1(x) = x, \\ p_{k+1}(x) &= \frac{2k+1}{k+1} x p_k(x) - \frac{k}{k+1} p_{k-1}(x), \quad k = 1, 2, \dots \end{aligned} \quad (2)$$

Additionally, the Legendre polynomials form an orthogonal set, such that

$$\int_{-1}^1 p_i(x) p_j(x) dx = h_i \delta_{ij}, \quad i, j = 0, 1, 2, \dots,$$

where  $\delta_{ij}$  is the Kronecker delta and  $h_i = 2/(2i+1)$  for any  $i = 0, 1, 2, \dots$ . According to approximation theory, a function  $f \in L^2([-1, 1])$  can be approximated by a series of Legendre polynomials basis functions in the following form

$$f_n(x) = \sum_{j=0}^n c_j p_j(x), \quad (3)$$

where  $c_j$  are unknown constants that need to be determined. Let us define

$$I(f_n, \lambda) = \oint_{-1}^1 \frac{f_n(x)}{\lambda - x} dx, \quad -1 < \lambda < 1. \quad (4)$$

From Eq. (3), we obtain

$$I(f_n, \lambda) = \sum_{j=0}^n c_j \oint_{-1}^1 \frac{p_j(x)}{\lambda - x} dx = \sum_{j=0}^n c_j q_j(\lambda),$$

where

$$q_0(\lambda) = \oint_{-1}^1 \frac{1}{\lambda - x} dx = \ln \left( \frac{1+\lambda}{1-\lambda} \right), \quad q_1(\lambda) = \lambda q_0(\lambda) - 2,$$

and

$$q_j(\lambda) = \oint_{-1}^1 \frac{p_j(x)}{\lambda - x} dx, \quad j = 2, 3, \dots$$

It can be easily shown that  $q_j(x)$  satisfies the recurrence relation given by Eq. (2), specifically

$$q_{j+1}(\lambda) = \frac{2j+1}{j+1} \lambda q_j(\lambda) - \frac{j}{j+1} q_{j-1}(\lambda), \quad j = 1, 2, \dots$$

From [13], we have derived a recurrence relation concerning certain values of  $f$  represented as follows

$$I(f_n, \lambda) = \sum_{i=0}^n \frac{q_{n+1}(\lambda) - q_{n+1}(t_i)}{(\lambda - t_i)p'_{n+1}(t_i)} f(t_i), \tag{5}$$

$$c_j = \sum_{i=0}^n \frac{f(t_i)}{h_j \mu_i} p_j(t_i), \quad j = 0, \dots, n, \tag{6}$$

Here,  $t_i$  are the roots of the equation  $p_{n+1}(x) = 0$  and  $\mu_i = \sum_{k=0}^n p_k^2(t_i)$  for each  $i = 0, 1, 2, \dots, n$ . Furthermore, several authors have studied Cauchy integral equations using various numerical methods. These methods include methods based on Bernstein polynomials [16, 17], the collocation method [5, 16], the Galerkin method [9], successive approximations [8], Chebyshev polynomial [12], the Adomian decomposition method [7], the differential transform method [1, 19], Gaussian elimination [11], Sinc approximations [2], GaussLegendre collocation [15], and so on.

The remainder of this paper is organized as follows. In Section 2, we describe the method for solving Cauchy-type singular integral equations. Error analysis and convergence of the method are discussed in Section 3. In Section 4, we demonstrate the efficiency of this method and present some numerical examples. Finally, Section 5 is devoted to the conclusion of this paper.

## 2 Algorithmic solution procedure

In this section, we approximate the unknown function  $f$  defined in Eq. (3) over the interval  $(-1, 1)$  using  $f_n$ , such that the following functional is minimized

$$\begin{aligned} R(f_n) &= \|g(\lambda) - f_n(\lambda) + \oint_{-1}^1 \frac{f_n(x)}{\lambda - x} dx\|_{L_\infty[-1,1]} \\ &= \|g(\lambda) - \sum_{j=0}^n c_j p_j(\lambda) + \sum_{j=0}^n c_j q_j(\lambda)\|_{L_\infty[-1,1]}, \quad -1 < \lambda < 1. \end{aligned} \tag{7}$$

Furthermore, by utilizing (5) we can express  $R(f_n)$  as follows

$$\begin{aligned} R(f_n) &= \|f_n(\lambda) - g(\lambda) - \sum_{i=0}^n \frac{q_{n+1}(\lambda) - q_{n+1}(t_i)}{(\lambda - t_i)p'_{n+1}(t_i)} f(t_i)\|_{L_\infty[-1,1]} \\ &= \|f_n(\lambda) - g(\lambda) - \sum_{i=0}^n V_i(\lambda) f(t_i)\|_{L_\infty[-1,1]}, \end{aligned}$$

where

$$V_i(\lambda) = \frac{q_{n+1}(\lambda) - q_{n+1}(t_i)}{(\lambda - t_i)p'_{n+1}(t_i)}.$$

From Eqs. (3), (6), and (7) we obtain

$$\begin{aligned} R(f_n) &= \left\| \sum_{j=0}^n \sum_{i=0}^n \frac{1}{\mu_i h_j} p_j(t_i) f(t_i) p_j(\lambda) - \sum_{i=0}^n f(t_i) V_i(\lambda) - g(\lambda) \right\|_{L_\infty[-1,1]} \\ &= \left\| \sum_{i=0}^n \left( \frac{1}{\mu_i} \sum_{j=0}^n \frac{1}{h_j} p_j(t_i) p_j(\lambda) - V_i(\lambda) \right) f(t_i) - g(\lambda) \right\|_{L_\infty[-1,1]}, \quad -1 < \lambda < 1, \end{aligned} \tag{8}$$

where  $f(t_i)$ ,  $i = 0, \dots, n$  are unknowns. Clearly, if  $g \in \pi_n$ , where  $\pi_n$  is the set of polynomials of degree at most  $n$ , then by selecting any  $n + 1$  arbitrary values from the interval  $(-1, 1)$ , we have  $R(f_n) = 0$ . Consequently, the minimization problem described in (8) has a unique solution. Additionally, if  $g \in L_2[-1, 1]$ , we can easily determine the minimum of  $R(f_n)$ .

In addition, if  $g \in C^{n+1}[-1, 1]$ , we can uniquely determine the unknown values  $f(t_i)$  for  $i = 0, \dots, n$  by interpolating  $g$  with a polynomial of degree at most  $n$ . To achieve this, we assume that

$$\lambda_m = -1 - \varepsilon + mk, \quad m = 0, 1, \dots, n. \quad (9)$$

where  $\varepsilon > 0$  is arbitrary and  $k = \frac{2-2\varepsilon}{n}$ . This choice ensures that the singular integral at the endpoint remains finite. By substituting these points into Eq. (8), we arrive at the conclusion

$$\sum_{j=0}^n c_j^* \left( p_j(\lambda_m) - \sum_{i=0}^n V_i(\lambda_m) p_j(t_i) \right) = g(\lambda_m), \quad m = 0, 1, \dots, n. \quad (10)$$

Due to the orthogonality of the basis  $\{p_n(x)\}$ , the system of equations in Eq. (10) possesses nonsingular properties, allowing us to obtain a unique solution for the unknowns. Consequently, by solving this system of  $n + 1$  equations with  $n + 1$  unknowns, we can compute the coefficients  $c_j^*$ . Finally, we define

$$f_n^*(x) = \sum_{j=0}^n c_j^* p_j(x). \quad (11)$$

The following algorithm demonstrate the step-by-step process of the proposed scheme through an algorithmic solution procedure.

#### Algorithm 1.

**Input:** Read  $n$  and  $\varepsilon > 0$ .

**Step 1:** Calculate  $k = (2 - 2\varepsilon)/n$  and  $\lambda_m = -1 - \varepsilon + mk$ , where  $m = 0, 1, \dots, n$ . Next, we consider the Legendre polynomials given in (2). The values  $t_i$ , which are the roots of the Legendre polynomials, are obtained by solving the equation  $p_{n+1}(x) = 0$ .

**Step 2:** Compute

$$\mu_i = \sum_{k=0}^n p_k^2(t_i), \text{ and } h_j = \frac{2}{2j+1}, \quad i, j = 0, 1, 2, \dots$$

**Step 3:** Calculate the recurrence relations given by the following equations

$$\begin{aligned} q_0(\lambda) &= \ln\left(\frac{1+\lambda}{1-\lambda}\right), & q_1(\lambda) &= \lambda q_0(\lambda) - 2, \\ q_{j+1}(\lambda) &= \frac{2j+1}{j+1} \lambda q_j(\lambda) - \frac{j}{j+1} q_{j-1}(\lambda), & j &= 1, 2, \dots \end{aligned}$$

**Step 4:** Solve Eq. (10) using any numerical method to find the solutions for the unknowns  $c_j^*$ .

**Output:** By solving the  $n + 1$  equations with  $n + 1$  unknowns, you will obtain the values of  $c_j^*$ . Finally, compute  $f_n^*(x)$  using (11).

### 3 Error analysis

we first establish the decay rates of the coefficients in the Legendre series expansion. We then demonstrate the error bounds of the truncated Legendre series in the uniform norm through two theorems. Finally, we derive the error analysis and convergence of the collocation method. We analyze the decay rates of the Legendre coefficients using the following theorem.

**Theorem 1.** *If  $f$  is analytic inside and on the Bernstein ellipse  $\varepsilon_\rho$  with foci at  $\pm 1$  and with the major and minor semi-axis summing to  $\rho > 1$ , then for each  $n \geq 0$ ,*

$$|f(x) - f_n(x)| \leq \frac{(2n\rho + 3\rho - 2n - 1)\ell(\varepsilon_\rho)M}{\pi\rho^{n+1}(\rho - 1)^2(1 - \rho^{-2})} = M_n, \quad x \in [-1, 1], \quad (12)$$

where  $M = \max_{z \in \varepsilon_\rho} |f(z)|$  and  $\ell(\varepsilon_\rho)$  denotes the length of the circumference of  $\varepsilon_\rho$ .

*Proof.* See [18] □

**Theorem 2.** *Assuming that the conditions of Theorem 1 are satisfied, we have  $f_n^* \rightarrow f$  as  $n \rightarrow \infty$ .*

*Proof.* To obtain the error estimation for the proposed approximation, let  $g_{p,n}$  be the interpolation polynomial of  $g$  at  $n + 1$  distinct nodes  $\lambda_m, m = 0, 1, \dots, n$ . Then, we get

$$\sum_{j=0}^n c_j^* p_j(\lambda) = g_{p,n}(\lambda) + \oint_{-1}^1 \frac{\sum_{j=0}^n c_j^* p_j(x)}{\lambda - x} dx.$$

Thus,

$$\sum_{j=0}^n c_j^* (p_j(\lambda) - q_j(\lambda)) = g_{p,n}(\lambda). \quad (13)$$

Now, let

$$\delta_n(\lambda) = f(\lambda) - f_n^*(\lambda) = f(\lambda) - \sum_{j=0}^n c_j^* p_j(\lambda).$$

Then, Eqs. (1) and (13) yield

$$\begin{aligned} \delta_n(\lambda) &= g(\lambda) + \oint_{-1}^1 \frac{f(x)}{\lambda - x} dx - \sum_{j=0}^n c_j^* p_j(\lambda) \\ &= \left( g(\lambda) + \oint_{-1}^1 \frac{f(x)}{\lambda - x} dx \right) - \left( g_{p,n}(\lambda) + \oint_{-1}^1 \frac{\sum_{j=0}^n c_j^* p_j(x)}{\lambda - x} dx \right) \\ &= (g(\lambda) - g_{p,n}(\lambda)) + \oint_{-1}^1 \frac{\delta_n(x)}{\lambda - x} dx. \end{aligned}$$

Consequently, we have

$$\begin{aligned} \left\| \delta_n(\lambda) - \oint_{-1}^1 \frac{\delta_n(x)}{\lambda - x} dx \right\|_\infty &= \|g(\lambda) - g_{p,n}(\lambda)\|_\infty \\ &\leq \frac{|g^{(n+1)}(\xi(\lambda))|}{(n+1)!} \prod_{i=0}^n |\lambda^2 - \lambda_i^2| \\ &= \frac{M_{n+1}}{(n+1)!}, \end{aligned}$$

where  $M_{n+1}$  is an upper bound for  $|g^{(n+1)}|$  in  $[-1, 1]$ . Clearly, we have

$$\lim_{n \rightarrow \infty} \left\| \delta_n(\lambda) - \oint_{-1}^1 \frac{\delta_n(x)}{\lambda - x} dx \right\|_{\infty} = 0.$$

The uniqueness theorem for solutions to equation (1) gives us

$$\lim_{n \rightarrow \infty} \left\| \delta_n(\lambda) \right\|_{\infty} = 0,$$

and thus, the proof is complete.  $\square$

## 4 Numerical examples

To illustrate the performance of our method, we dedicate this section to presenting numerical results through several examples. In these computations, each table displays the absolute errors of our approximate solutions.


**Example 1.** Consider the integral equation

$$f(\lambda) = \frac{\lambda + 2Ln3}{\lambda + 2} - \frac{\lambda}{\lambda + 2} Ln \left| \frac{\lambda + 1}{\lambda - 1} \right| + \int_{-1}^1 \frac{f(x)}{\lambda - x} dx,$$

where the exact solution is given by  $f(\lambda) = \frac{\lambda}{\lambda + 2}$ .

**Table 1:** Absolute errors for Example 1.

$N$	$\varepsilon = 0.001$	$\varepsilon = 0.0001$
6	2.289E-3	1.252E-3
8	3.704E-4	1.759E-4
10	6.241E-5	2.513E-5
12	1.098E-5	3.625E-6



**Figure 1:** Comparison between the exact and approximate solutions of Example 1 with  $N = 12$ .

**Example 2.** Consider the integral equation as described in [3]

$$k\varphi = \varphi(t) + \int_L \frac{\varphi(\tau)}{\tau - t} dt = f(t),$$


where

$$f(t) = \frac{1}{1+t^2} \left( t^3 \text{Ln}\left(\frac{1-t}{1+t}\right) + t^3 + 2t^2 + \frac{4-\pi}{2} \right), \quad k(t, \tau) = 0,$$


and  $L = [-1, 1]$ . The exact solution is given by  $\varphi(t) = \frac{t^3}{1+t^2}$ .

**Table 2:** Comparison of absolute errors [3] with the present method considering  $\varepsilon = 0.0001$ .

$N$	Present method	Method used in [3]
4	1.488E-1	1.129E-1
8	1.162E-2	3.338E-2
16	2.710E-4	4.933E-3



**Figure 2:** Comparison between the exact and approximate solutions of Example 2 with  $N = 16$



**Figure 3:** Absolute errors for Example 2 with different values of  $N$  considering  $\varepsilon = 0.0001$ .

**Example 3.** Consider the integral equation from [15]

$$\lambda x(s) + \mu \int_{-1}^1 \frac{x(t)}{t-s} dt = \xi(s), \quad s \in (-1, 1),$$


where  $\lambda = \mu = 1$ , and

$$\begin{aligned} \xi(s) = & \sinh(s) - \operatorname{chi}(-s-1) \sinh(s) + \operatorname{chi}(1-s) \sinh(s) + \operatorname{Ln}(-1-s) \sinh(s) - \operatorname{Ln}(1-s) \sinh(s) \\ & + \operatorname{Ln}\left(\frac{1-s}{1+s}\right) \sinh(s) + \operatorname{cosh}(s) \operatorname{shi}(1-s) + \operatorname{cosh}(s) \operatorname{shi}(1+s). \end{aligned}$$

In this context, *chi* and *shi* refer to the hyperbolic cosine integral and hyperbolic sine integral, respectively. The exact solution is given by  $x(s) = \sinh(s)$ .


**Table 3:** Comparison of absolute errors between the present method considering  $\varepsilon = 0.0001$  and the method in [15].

Node	Presented method			Method of [15]		
	$N = 3$	$N = 5$	$N = 7$	$N = 3$	$N = 5$	$N = 7$
-1.0	$2.21e-4$	$3.29e-6$	$2.34e-8$	$1.91e-3$	$1.81e-5$	$8.90e-8$
-0.8	$6.80e-4$	$5.97e-7$	$2.89e-8$	$3.27e-4$	$2.71e-6$	$1.03e-8$
-0.6	$2.80e-4$	$1.27e-6$	$1.56e-9$	$4.56e-4$	$3.09e-6$	$2.75e-10$
-0.4	$1.81e-4$	$2.89e-6$	$1.61e-8$	$7.13e-4$	$4.86e-7$	$4.69e-9$
-0.2	$8.84e-5$	$3.93e-6$	$6.74e-9$	$5.31e-4$	$1.50e-6$	$1.59e-8$
-0.2	$4.31e-4$	$1.15e-6$	$1.13e-9$	$7.62e-6$	$2.35e-6$	$5.44e-10$
0.0	$7.10e-4$	$6.16e-7$	$1.28e-8$	$4.36e-4$	$5.16e-6$	$7.95e-9$
0.4	$7.88e-4$	$2.84e-6$	$3.03e-9$	$3.80e-4$	$1.85e-6$	$1.33e-8$
0.6	$2.21e-4$	$6.49e-7$	$1.93e-8$	$2.72e-4$	$2.78e-6$	$1.17e-8$
0.8	$5.63e-4$	$3.64e-7$	$2.43e-9$	$9.57e-4$	$1.95e-6$	$7.49e-9$
1.0	$8.19e-5$	$5.14e-7$	$2.23e-10$	$8.32e-5$	$2.08e-6$	$1.23e-8$



**Figure 4:** Comparison between the exact and approximate solutions for Example 3 with  $N = 7$ .





**Figure 5:** Absolute errors for Example 3 in [15] (left) and the absolute errors of the solutions obtained in this paper considering  $\varepsilon = 0.0001$  (right).

This study employs Legendre orthogonal polynomial expansions, offering a robust framework for approximating solutions due to their inherent orthogonality. This approach significantly reduces the computational effort compared to methods that utilize Bessel basis polynomials and Gauss-Legendre collocation points, making it a more efficient option for solving these equations.

## 5 Conclusion

We have used an expansion approach for the unknown function and the Cauchy kernel in this study. The unknown coefficients were defined by applying the collocation method. Using a system of linear equations, we can compute an approximate solution and eliminate singularities with this method. Through the provided examples, we show that, although our numerical solution is straightforward and quick, it is also very accurate and efficient when compared to other approaches used in related studies.

## References

- [1] M. Abdulkawi, *Solution of Cauchy type singular integral equations of the first kind by using differential transform method*, Appl. Math. Model. **39** (2015) 2107–2118.
- [2] M.S. Akel, H.S. Hussein, *Numerical treatment of solving singular integral equations by using sinc approximations*, Appl. Math. Comput. **218** (2011) 3565–3573.
- [3] A. Chakrabarti, G. Vanden Berghe, *Approximate solution of singular integral equations*, Appl. Math. Lett. **17** (2004) 553–559.
- [4] D. Elliott, *Rates of convergence for the method of classical collocation for solving integral equations*, SIAM J. Numer. Anal. **21** (1984) 136–148.
- [5] E.G. Ladopoulos, *Singular Integral Equations, Linear and Non-linear Theory and its Applications in Science and Engineering*, Springer-Verlag, Berlin, New York, 2000.
- [6] A. Linkov, *Boundary Integral Equations in Elasticity Theory*, Kluwer, Dordrecht, 2002.
- [7] B.N. Mandal, G. Bera, *Approximate solution of a class of singular integral equations of second kind*, J. Comput. Appl. Math. **206** (2007) 189–195.

- [8] A. Mennouni, S. Guedjiba, *A note on solving Cauchy integral equations of the first kind by iterations*, Appl. Math. Comput. **217** (2011) 7442–7447.
- [9] A. Mennouni, *Piecewise constant Galerkin method for a class of Cauchy singular integral equations of the second kind in  $L^2$* , Appl. Math. Comput. **326** (2017) 268–272.
- [10] N.I. Muskhelishvili, *Singular Integral Equations*, Groningen Noordhoff, Holland, 1953.
- [11] H.X. Mamatova, Z.K. Eshkuvatov, S. Ismail, *A Hybrid Method for All Types of Solutions of the System of Cauchy-Type Singular Integral Equations of the First Kind*, Mathematics **11** (2023) 4404.
- [12] V.N. Nfor, P.L. Ndikum, *Solving singular integral equations of the second kind using Chebyshev polynomials*, Afr. J. Math. Comput. Sci. Res. **16** (2023) 27–35.
- [13] D.F. Paget, D. Elliott, *An algorithm for the numerical evaluation of certain Cauchy principal value integrals*, Numer. Math. **19** (1972) 373–385.
- [14] A.D. Polyanin, A.V. Manzhirov, *Handbook of Integral Equations*, Second Edition, Front Cover CRC Press, Math, 2008.
- [15] A. Seifi, *Numerical solution of certain Cauchy singular integral equations using a collocation scheme*, Adv. Differ. Equ. **2020** (2020) 1–15.
- [16] A. Seifi, T. Lotfi, T. Allahviranloo, M. Paripour, *An effective collocation technique to solve the singular Fredholm integral equations with Cauchy kernel*, Adv. Differ. Equ. **280** (2017) 1-18.
- [17] V. Sharma, A. Setia, R.P. Agarwal, *Numerical solution for system of Cauchy type singular integral equations with its error analysis in complex plane*, Appl. Math. Comput, **328** (2018) 338–352.
- [18] H. Wang, S. Xiang, *On the convergence rates of Legendre approximation*, Math. Comput. J **81** (2012) 861–877.
- [19] L. Yang, Z. Chen, K. Xie, *An efficient method for approximate solution of a singular integral equation with Cauchy kernel*, J. Comput. Appl. Math. **352** (2019) 50–61.